

On the quasi-component of pseudocompact abelian groups *

D. Dikranjan¹ and Gábor Lukács²

February 20, 2012

Abstract

In this paper, we describe the relationship between the quasi-component $q(G)$ of a (perfectly) minimal pseudocompact abelian group G and the quasi-component $q(\tilde{G})$ of its completion. Specifically, we characterize the pairs (C, A) of compact connected abelian groups C and subgroups A such that $A \cong q(G)$ and $C \cong q(\tilde{G})$. As a consequence, we show that for every positive integer n or $n = \omega$, there exist plenty of abelian pseudocompact perfectly minimal n -dimensional groups G such that the quasi-component of G is not dense in the quasi-component of the completion of G .

1. Introduction

A Tychonoff space X is *pseudocompact* if every continuous real-valued map on X is bounded. If a topological group G is pseudocompact, then its completion \tilde{G} is compact, that is, G is *precompact* (cf. [4, 1.1]), which allows for the following characterization of pseudocompact groups.

Theorem 1.1. ([4]) *A topological group G is pseudocompact if and only if G is precompact and G_δ -dense in \tilde{G} , in which case $\tilde{G} = \beta G$.*

A Tychonoff space is *zero-dimensional* if it has a base consisting of *clopen* (open-and-closed) sets. For a topological group G , let G_0 and $q(G)$ denote the connected component and the quasi-component of the identity, respectively. For every group G , the quotient G/G_0 is *hereditarily disconnected* (i.e., $(G/G_0)_0$ is trivial), and $G/q(G)$ is *totally disconnected* (i.e., $q(G/q(G))$ is trivial). While in the class of locally compact groups (or spaces) these notions of disconnectedness coincide with zero-dimensionality, this is not the case in general. Indeed, both G/G_0 and $G/q(G)$ may fail to be zero-dimensional, or to admit a coarser zero-dimensional group topology, even in the presence of additional compactness-like properties (cf. [3, 7.7], [17], [8, Theorem D]). There is a close relationship between connectedness and disconnectedness properties of pseudocompact groups and those of their completions, which is summarized in the next theorem.

* 2010 Mathematics Subject Classification: Primary 22A05, 54D25, 54H11; Secondary 22C05, 54D05, 54D30

¹ The first author acknowledges the financial aid received from grant MTM2009-14409-C02-01.

² The second author gratefully acknowledges the generous financial support received from NSERC, which enabled him to do this research.

Theorem 1.2. *Let G be a pseudocompact group. Then:*

- (a) ([5, 1.4]) $q(G) = q(\tilde{G}) \cap G = (\tilde{G})_0 \cap G$;
- (b) ([26]) G is zero-dimensional if and only if \tilde{G} is zero-dimensional;
- (c) ([7, 1.7]) $G/q(G)$ is zero-dimensional if and only if $q(G)$ is dense in $(\tilde{G})_0 = q(\tilde{G})$;
- (d) ([2, 4.11(b)]) G/G_0 is zero-dimensional if and only if G_0 is dense in $(\tilde{G})_0 = q(\tilde{G})$, in which case $G_0 = q(G)$.

We say that a pair (C, A) of a group C and its subgroup A is *realized* by a group G if $A \cong q(G)$ and $C \cong q(\tilde{G})$. In a previous paper, the authors obtained sufficient conditions for a pair to be realizable by pseudocompact abelian groups that satisfy various degrees of so-called minimality properties (cf. [8, Theorem D']). In this paper, we obtain a complete characterization (i.e., necessary and sufficient conditions) for the same.

Recall that a (Hausdorff) topological group G is *minimal* if there is no coarser (Hausdorff) group topology on G (cf. [23] and [11]). For a pseudocompact group G , the quotient $G/q(G)$ is totally disconnected and pseudocompact, and thus, by an unpublished result of Shakhmatov, admits a coarser zero-dimensional group topology (cf. [8, Theorem B]). Therefore, Shakhmatov's result implies that for a pseudocompact group G , if $G/q(G)$ is minimal, then it is zero-dimensional. If in addition G is minimal and abelian, then the converse is also true.

Theorem 1.3. ([6, Theorem 3], [7, 1.7]) *Let G be a minimal pseudocompact abelian group. Then the following statements are equivalent:*

- (i) $G/q(G)$ is zero-dimensional;
- (ii) $q(G)$ is dense in $(\tilde{G})_0 = q(\tilde{G})$;
- (iii) $G/q(G)$ is minimal.

A group G is *totally minimal* if every (Hausdorff) quotient of G is totally minimal (cf. [9]); G is *perfectly (totally) minimal* if the product $G \times H$ is (totally) minimal for every (totally) minimal group H (cf. [25]). In an earlier paper, the authors proved the following:

Theorem 1.4. ([8, Theorem D']) *Let C be a connected compact abelian group, and A a subgroup of C . Then there exists a pseudocompact abelian group G such that $A \cong q(G)$ and $C \cong (\tilde{G})_0 = q(\tilde{G})$, and in particular, $\dim G = \dim C$. Furthermore, if A is dense in C and*

- (a) A is minimal, then G may be chosen to be minimal;
- (b) A is totally minimal, then G may be chosen to be totally minimal;
- (c) A is perfectly minimal, then G may be chosen to be perfectly minimal;
- (d) A is perfectly totally minimal, then G may be chosen to be perfectly totally minimal.

By Theorem 1.3, if G is a totally minimal pseudocompact abelian group, then $q(G)$ is dense in $(\tilde{G})_0 = q(\tilde{G})$. Thus, the condition that A is dense in C is not only sufficient, but also necessary in parts (b) and (d) of Theorem 1.4.

In this paper, we show that in order to realize (C, A) by a (perfectly) minimal pseudocompact abelian group, A need not be dense in C , but a milder condition is both sufficient and necessary. A subgroup E of a topological group G is *essential*, and we put $E \leq_e G$, if for every non-trivial closed normal subgroup N of G , the intersection $E \cap N$ is non-trivial.

Theorem A. *Let C be a compact connected abelian group, and A a subgroup. Then:*

- (a) *(C, A) can be realized by a minimal pseudocompact abelian group if and only if $A \leq_e C$;*
- (b) *(C, A) can be realized by a perfectly minimal pseudocompact abelian group if and only if A is perfectly minimal and $A \leq_e C$;*
- (c) *(C, A) can be realized by a (perfectly) totally minimal pseudocompact abelian group if and only if A is (perfectly) totally minimal and dense in C .*

Remark 1.5. If A is an essential subgroup of a compact group C , then in particular, it is essential in its completion $\tilde{A} \subseteq C$, and thus A is minimal (see Lemma 2.4(a) and Theorem 2.1). Therefore, the condition $A \leq_e C$ in Theorem A(a) implies that A is minimal.

By applying Theorem A to the character group $\text{hom}(\mathbb{Q}, \mathbb{R}/\mathbb{Z})$ equipped with the topology of pointwise convergence and the annihilator of \mathbb{Z} in it, we also obtain a family of “pathological” examples.

Theorem B. *For every positive integer n or $n = \omega$, there exists an abelian pseudocompact perfectly minimal group G_n with $\dim G_n = n$ such that $q(G_n)$ is not dense in $q(\tilde{G}_n) = (\tilde{G}_n)_0$, or equivalently, $G_n/q(G_n)$ is not minimal.*

The proofs of Theorems A and B are presented in §3; they are based on preservation properties of essential subgroups established in §2.

2. Preliminaries: Essential and minimal subgroups

While preservation of minimality under formation of closed subgroups and products has been thoroughly studied (cf. [20, Proposition 2.3, Lemma 3.1], [11, Théorème 1-2], [24, 9], and [12, (6), (3)]), it appears that preservation of essentiality has not been well investigated in the context of topological groups. Our aim in this section is to remedy this state of affairs in the realm of abelian groups.

The relationship between minimality and essential subgroups was discovered independently by Stephenson and Prodanov, and generalized by Banaschewski (cf. [23, Theorem 2], [19], and [1, Propositions 1 and 2]).

Theorem 2.1. ([10, 2.5.1], [16, 3.21]) *Let G be a topological group, and D a dense subgroup. Then D is minimal if and only if G is minimal and $D \leq_e G$.*

The celebrated Prodanov-Stoyanov Theorem states that every minimal abelian group is pre-compact (cf. [21] and [22]), and allows for a complete characterization of minimality of abelian groups using the notion of an essential subgroup.

Theorem 2.2. ([10, 2.5.2], [16, 3.31]) *An abelian topological group G is minimal if and only if its completion \tilde{G} is compact and $G \leq_e \tilde{G}$.*

The next two easy lemmas, whose proofs have been omitted, describe elementary properties of the essentiality relation.

Lemma 2.3. *Let G be a topological group, and E and H subgroups of G such that $E \subseteq H$.*

(a) *If $E \leq_e H$ and $H \leq_e G$, then $E \leq_e G$.*

(b) *If H is dense in G , then $E \leq_e G$ if and only if $E \leq_e H$ and $H \leq_e G$.* □

Lemma 2.4. *Let G be an abelian topological group, and E and H subgroups.*

(a) *If $E \subseteq H$, then $E \leq_e G$ if and only if $E \leq_e H$ and $H \leq_e G$.*

(b) *If H is closed in G and $E \leq_e G$, then $E \cap H \leq_e H$.* □

Remark 2.5. Every closed subgroup of a minimal abelian group is minimal (cf. [20, Proposition 2.3]). Indeed, let G be a minimal abelian group, and M a closed subgroup. Then, by Theorem 2.2, \tilde{G} is compact and $G \leq_e \tilde{G}$. So, by Lemma 2.4(b), $M = G \cap (\text{cl}_{\tilde{G}} M) \leq_e \text{cl}_{\tilde{G}} M = \tilde{M}$. Thus, by Theorem 2.2, M is minimal. (In fact, every closed central subgroup of a minimal group is minimal; cf. [10, 7.2.5] and [16, 3.26].) It follows that every closed subgroup of a totally minimal abelian group is totally minimal, and in fact, every closed central subgroup of a totally minimal group is totally minimal (cf. [20, Lemma 3.1], [10, 7.2.5], and [16, 3.27]).

We turn now to preservation of essentiality under formation of products. A topological group G is *elementwise compact* if every $g \in G$ is contained in a compact subgroup of G , or equivalently, $\text{cl}_G \langle g \rangle$ is compact for every $g \in G$ (cf. [15, 5.4]). Stephenson showed that every elementwise compact minimal group is perfectly minimal (cf. [24, 9]). The next lemma is a natural extension of Stephenson's result.

Lemma 2.6. *Let G be a topological group, and $E \leq_e G$. If L is an elementwise compact group, then $L \times E \leq_e L \times G$.*

PROOF. Let $\pi_2: L \times G \rightarrow G$ denote the canonical projection, and let N be a non-trivial closed normal subgroup of $L \times G$. If $\pi_2(N)$ is trivial, then $N \subseteq L \times \{e\}$, and consequently $N \cap (L \times E)$ is non-trivial. Thus, we may assume that $\pi_2(N)$ is non-trivial. We prove the statement in two steps.

Step 1. Suppose that L is compact. Then π_2 is a closed map (cf. [13, 3.1.16]), and so $\pi_2(N)$ is a closed normal subgroup of G . Thus, $\pi_2(N) \cap E$ is non-trivial, because $E \leq_e G$. Therefore, the intersection $N \cap (L \times E)$ is non-trivial, as required.

Step 2. In the general case, let $x = (l, g) \in N$ be such that $g \neq e$. Since L is elementwise compact, there is a compact subgroup S of L such that $l \in S$. Put $N' := N \cap (S \times G)$. Then N' is a closed normal subgroup of $S \times G$, and it is non-trivial, because $x \in N'$. By what we have shown so far, $N' \cap (S \times E)$ is non-trivial, and in particular, $N \cap (L \times E)$ is non-trivial, as desired. □

The next example shows that elementwise compactness cannot be replaced with precompactness, minimality, or completeness in Lemma 2.6.

Example 2.7. Let p be a prime, \mathbb{Z}_p the group of p -adic integers, and (\mathbb{Z}, τ_p) the integers equipped with the p -adic topology. Since (\mathbb{Z}, τ_p) is a minimal group whose completion is \mathbb{Z}_p (cf. [10, 2.5.6]), by Theorem 2.2, $\mathbb{Z} \leq_e \mathbb{Z}_p$. However, $\mathbb{Z} \times \mathbb{Z}$ is not essential in $(\mathbb{Z}, \tau_p) \times \mathbb{Z}_p$ or $\mathbb{Z} \times \mathbb{Z}_p$ (in the latter, the first component is equipped with the discrete topology). Indeed, if $\xi \in \mathbb{Z}_p \setminus \mathbb{Z}$, then $F := \langle (1, \xi) \rangle$ is a non-trivial closed subgroup of $(\mathbb{Z}, \tau_p) \times \mathbb{Z}_p$ (and thus of $\mathbb{Z} \times \mathbb{Z}_p$) such that $F \cap (\mathbb{Z} \times \mathbb{Z})$ is trivial.

It is also worth noting that preservation of essentiality does not imply precompactness, minimality, nor completeness, as the next example demonstrates.

Example 2.8. Let p be a prime. Let L_1 denote the direct sum $\mathbb{Z}_p^{(\omega)}$ equipped with the subgroup topology induced by the direct product \mathbb{Z}_p^ω , let L_2 denote the direct sum $(\mathbb{R}/\mathbb{Z})^{(\omega)}$ equipped with the box topology, and put $L := L_1 \times L_2$. Then L is elementwise compact, because elementwise compactness is preserved under formation of products, coproducts, sums, and Σ -products. Thus, by Lemma 2.6, L preserves essentiality, that is, $E \leq_e G$ implies $L \times E \leq_e L \times G$. Nevertheless, L is neither complete nor precompact, and in particular, by Theorem 2.2, L is not minimal.

These examples provide a natural motivation for the following problem.

Problem 2.9. Characterize the topological groups L with the property that

- (a) for every topological group G , if $E \leq_e G$, then $L \times E \leq_e L \times G$;
- (b) for every abelian topological group G , if $E \leq_e G$, then $L \times E \leq_e L \times G$;
- (c) for every topological group G , if $E \leq_e G$ and E is minimal, then $L \times E \leq_e L \times G$.

We provide an answer to Problem 2.9(c) in the special case where L is minimal and abelian.

Lemma 2.10. Let M be a minimal abelian group, G a topological group, and $E \leq_e G$. If $M \times E$ is minimal, then:

- (a) $M \times E \leq_e \widetilde{M} \times E$;
- (b) $M \times E \leq_e \widetilde{M} \times G$;
- (c) $M \times E \leq_e M \times G$.

PROOF. (a) By Theorem 2.1, $M \times E \leq_e \widetilde{M} \times \widetilde{E}$, because $M \times E$ is minimal. Consequently, by Lemma 2.3(b), $M \times E \leq_e \widetilde{M} \times E$, since $\widetilde{M} \times E$ is dense in $\widetilde{M} \times \widetilde{E}$.

(b) By Theorem 2.2, \widetilde{M} is compact, and so, by Lemma 2.6, $\widetilde{M} \times E \leq_e \widetilde{M} \times G$. By Lemma 2.3(a), combining this with what has been shown in part (a) yields $M \times E \leq_e \widetilde{M} \times G$.

(c) As $M \times G$ is dense in $\widetilde{M} \times G$, by Lemma 2.3(b), part (b) implies that $M \times E \leq_e M \times G$. \square

Theorem 2.11. Let M be a minimal abelian group. The following statements are equivalent:

- (i) M is perfectly minimal;
- (ii) for every topological group G , if $E \leq_e G$ and E is minimal, then $M \times E \leq_e M \times G$;
- (iii) for every abelian topological group G , if $E \leq_e G$ and E is minimal, then $M \times E \leq_e M \times G$;
- (iv) for every minimal abelian group M' , the product $M \times M'$ is minimal.

PROOF. The implication (i) \Rightarrow (ii) follows by Lemma 2.10(c), and the implication (ii) \Rightarrow (iii) is trivial. The equivalence (i) \Leftrightarrow (iv) is an immediate consequence of the following theorem of Stoyanov: A topological group G is perfectly minimal if and only if $G \times (\mathbb{Z}, \tau_p)$ is minimal for every prime p (cf. [25]). We turn now to the remaining implication.

(iii) \Rightarrow (iv): Let M' be a minimal abelian group. By Theorem 2.2, \widetilde{M} and \widetilde{M}' are compact, and $M \leq_e \widetilde{M}$ and $M' \leq_e \widetilde{M}'$. So, by (iii), $M \times M' \leq_e M \times \widetilde{M}'$, and by Lemma 2.6, $M \times \widetilde{M}' \leq_e \widetilde{M} \times \widetilde{M}'$. Therefore, by Lemma 2.3(a), $M \times M' \leq_e \widetilde{M} \times \widetilde{M}'$. Hence, by Theorem 2.2, $M \times M'$ is minimal. \square

Remark 2.12. It follows from Remark 2.5 and Theorem 2.11 that every closed subgroup of a perfectly minimal abelian group is perfectly minimal.

3. Proof of Theorems A and B

We prove Theorem A by establishing the following more elaborate statement.

Theorem A'. *Let A be an essential subgroup of a connected compact abelian group C . Then there exists an abelian group G such that*

- (a) G is pseudocompact;
- (b) $A \cong q(G)$;
- (c) $C \cong q(\tilde{G}) = (\tilde{G})_0$, and in particular, $\dim G = \dim C$;
- (d) G is minimal.

Furthermore, if A is perfectly minimal, then G may be chosen to be perfectly minimal.

We first show how Theorem A follows from Theorem A'.

PROOF OF THEOREM A. Sufficiency of the conditions in (a) and (b) follows from Theorem A', while sufficiency of (c) was already shown in [8, Theorem D']. Thus, we may turn to the necessity of the conditions.

(a) Let G be a minimal pseudocompact abelian group. Then, by Theorem 2.2, $G \leq_e \tilde{G}$, and thus, by Lemma 2.4, $G \cap q(\tilde{G}) \leq_e q(\tilde{G})$. By Theorem 1.2, $q(G) = q(\tilde{G}) \cap G$, and therefore $q(G) \leq_e q(\tilde{G})$.

(b) If G is a perfectly minimal pseudocompact abelian group, then by Remark 2.12, $q(G)$ is also perfectly minimal, and by part (a), $q(G) \leq_e q(\tilde{G})$.

(c) Let G be a (perfectly) totally minimal pseudocompact abelian group. Then $G/q(G)$ is minimal, and by Theorem 1.3, $q(G)$ is dense in $q(\tilde{G})$. It follows from Remarks 2.5 and 2.12 that $q(G)$ is (perfectly) totally minimal, being a closed subgroup of a (perfectly) totally minimal abelian group. \square

We turn now to the proof of Theorem A', and to that end, we recall two technical lemmas.

Lemma 3.1. ([8, 5.2]) *For every infinite cardinal λ , there exists a pseudocompact zero-dimensional group H such that:*

- (i) H is perfectly totally minimal;
- (ii) $r_0(\tilde{H}/H) \geq 2^\lambda$.

Lemma 3.2. ([8, 5.3]) *Let K_1 and K_2 be compact topological groups, and let $h: K_1 \rightarrow K_2$ be a surjective homomorphism such that $\ker h$ is G_δ -dense in K_1 . Then the graph Γ_h of h is a G_δ -dense subgroup of the product $K_1 \times K_2$, and in particular, Γ_h is pseudocompact.*

PROOF OF THEOREM A'. Put $\lambda = w(C)$, and let H be the group provided by Lemma 3.1. Since $r_0(\tilde{H}/H) \geq 2^\lambda$, the quotient \tilde{H}/H contains a free abelian group F of rank 2^λ . As $|C| \leq 2^\lambda$, one may pick a surjective homomorphism $h_1: F \rightarrow C$. The group C is divisible, because it is compact and connected (cf. [14, 24.25]). Thus, h_1 can be extended to a surjective homomorphism $h_2: \tilde{H}/H \rightarrow C$.

Let $h: \tilde{H} \rightarrow C$ denote the composition of h_2 with the canonical projection $\tilde{H} \rightarrow \tilde{H}/H$. By Theorem 1.1, H is G_δ -dense in \tilde{H} , because H is pseudocompact. Thus, $\ker h$ is G_δ -dense in \tilde{H} ,

because $H \subseteq \ker h$. Clearly, h is surjective. Therefore, by Lemma 3.2, the graph Γ_h of h is G_δ -dense in the product $\tilde{H} \times C$.

Put $G := \Gamma_h + (\{0\} \times A)$. Since Γ_h is G_δ -dense in $\tilde{H} \times C$ and contained in G , the group G is G_δ -dense too. Thus, $\tilde{G} = \tilde{H} \times C$, and by Theorem 1.1, G is pseudocompact. As H is zero-dimensional, $q(\tilde{G}) = (\tilde{G})_0 = \{0\} \times C$, and by Theorem 1.2(a), $q(G) = q(\tilde{G}) \cap G = \{0\} \times A$.

We check now that $\dim G = \dim C$. Since G is pseudocompact, by Theorem 1.1, $\tilde{G} = \beta G$, and so $\dim G = \dim \beta G = \dim \tilde{G}$ (cf. [13, 7.1.17]). As H is zero-dimensional and pseudocompact, by Theorem 1.2(b), $\dim \tilde{H} = 0$. Thus, by Yamanoshita's Theorem, $\dim \tilde{G} = \dim \tilde{H} + \dim C = \dim C$ (cf. [27], [18, Corollary 2], and [10, 3.3.12]). Therefore, $\dim G = \dim C$.

We turn now to minimality properties of G . The group G always contains the product $H \times A$. Since C is compact, so is $\tilde{A} = \text{cl}_C A$, and by Lemma 2.4(a), $A \leq_e \tilde{A}$, because $A \leq_e C$. Thus, by Theorem 2.2, A is minimal, and consequently, $H \times A$ is minimal, as H is perfectly minimal. Therefore, by Lemma 2.10(b), $H \times A \leq_e \tilde{H} \times C = \tilde{G}$. Hence, by Lemma 2.4(a), $G \leq_e \tilde{G}$, and by Theorem 2.2, G is minimal.

Suppose now that A is perfectly minimal, and let M' be a minimal abelian group. Then $H \times A$ is perfectly minimal, and so $H \times A \times M'$ is minimal. Thus, by Lemma 2.10(b), $H \times A \times M' \leq_e \tilde{G} \times \tilde{M}'$. Therefore, by Lemma 2.4(a), $G \times M' \leq_e \tilde{G} \times \tilde{M}'$, because $G \times M'$ contains $H \times A \times M'$. Hence, by Theorem 2.2, $G \times M'$ is minimal. Consequently, by Theorem 2.11, G is perfectly minimal, as desired. \square

In preparation for the proof of Theorem B, we recall some remarkable properties of the Pontryagin dual of the discrete group of the rationals. Let $\mathbb{T} := \mathbb{R}/\mathbb{Z}$, and put $\hat{\mathbb{Q}} := \text{hom}(\mathbb{Q}, \mathbb{T})$, equipped with the topology of pointwise convergence. The group $\hat{\mathbb{Q}}$ is compact, and it is connected, because \mathbb{Q} is torsion-free (cf. [10, 3.3.8]).

Lemma 3.3. ([10, 3.6.2, 3.6.5]) *Let $\mathbb{Z}^\perp := \{\chi \in \hat{\mathbb{Q}} \mid \chi(\mathbb{Z}) = 0\}$ denote the annihilator of \mathbb{Z} in $\hat{\mathbb{Q}}$.*

- (a) \mathbb{Z}^\perp is a compact essential subgroup of $\hat{\mathbb{Q}}$;
- (b) $\mathbb{Z}^\perp \cong \prod_{p \in \mathbb{P}} \mathbb{Z}_p$, and in particular, $\dim \mathbb{Z}^\perp = 0$;
- (c) $\hat{\mathbb{Q}}/\mathbb{Z}^\perp \cong \mathbb{T}$, and in particular, $\dim \hat{\mathbb{Q}} = 1$.

PROOF OF THEOREM B. Suppose that $n < \omega$. Put $C_n := (\hat{\mathbb{Q}})^n$ and $A_n := (\mathbb{Z}^\perp)^n$. By Lemma 3.3, \mathbb{Z}^\perp is a compact essential subgroup of $\hat{\mathbb{Q}}$. Thus, by applying Lemmas 2.6 and 2.3(a) repeatedly, one obtains that A_n is a compact essential subgroup of C_n . It follows from Lemma 3.3 and Yamanoshita's Theorem that $\dim C_n = n$ (cf. [27], [18, Corollary 2], and [10, 3.3.12]). By Theorem A', there exists a perfectly minimal pseudocompact group G_n such that $A_n \cong q(G_n)$, $C_n \cong q(\tilde{G}_n)$, and $\dim G_n = \dim C_n = n$. Clearly, $q(G_n)$ is not only not dense, but in fact it is closed in $q(\tilde{G}_n)$.

For $n = \omega$, put $G_\omega = G_1 \times \mathbb{T}^\omega$. Then $\dim G_\omega \geq \omega$, and $q(G_\omega) = q(G_1) \times \mathbb{T}^\omega \cong \mathbb{Z}^\perp \times \mathbb{T}^\omega$, while

$$q(\tilde{G}_\omega) = (\tilde{G}_\omega)_0 = (\tilde{G}_1 \times \mathbb{T}^\omega)_0 = (\tilde{G}_1)_0 \times \mathbb{T}^\omega \cong \hat{\mathbb{Q}} \times \mathbb{T}^\omega.$$

Thus, $q(G_\omega)$ is not dense in $q(\tilde{G}_\omega)$, as required. The last statement follows from Theorem 1.3. \square

Acknowledgements

We are grateful to Karen Kipper for her kind help in proofreading this paper for grammar and punctuation. We wish to thank the anonymous referee for the constructive comments that led to an improved presentation of this paper.

References

- [1] B. Banaschewski. Minimal topological algebras. *Math. Ann.*, 211:107–114, 1974.
- [2] W. W. Comfort and G. Lukács. Locally precompact groups: (local) realcompactness and connectedness. *J. Lie Theory*, 20:347–374, 2010.
- [3] W. W. Comfort and J. van Mill. Concerning connected, pseudocompact abelian groups. *Topology Appl.*, 33(1):21–45, 1989.
- [4] W. W. Comfort and K. A. Ross. Pseudocompactness and uniform continuity in topological groups. *Pacific J. Math.*, 16:483–496, 1966.
- [5] D. Dikranjan. Connectedness and disconnectedness in pseudocompact groups. *Rend. Accad. Naz. Sci. XL Mem. Mat. (5)*, 16:211–221, 1992.
- [6] D. Dikranjan. Dimension and connectedness in pseudo-compact groups. *C. R. Acad. Sci. Paris Sér. I Math.*, 316(4):309–314, 1993.
- [7] D. Dikranjan. Zero-dimensionality of some pseudocompact groups. *Proc. Amer. Math. Soc.*, 120(4):1299–1308, 1994.
- [8] D. Dikranjan and G. Lukács. On zero-dimensionality and the connected component of locally pseudocompact groups. *Proc. Amer. Math. Soc.*, 139(8):2995–3008, 2011.
- [9] D. Dikranjan and I. R. Prodanov. Totally minimal topological groups. *Annuaire Univ. Sofia Fac. Math. Méc.*, 69:5–11 (1979), 1974/75.
- [10] D. Dikranjan, I. R. Prodanov, and L. N. Stoyanov. *Topological groups*, volume 130 of *Monographs and Textbooks in Pure and Applied Mathematics*. Marcel Dekker Inc., New York, 1990. Characters, dualities and minimal group topologies.
- [11] D. Doitchinov. Produits de groupes topologiques minimaux. *Bull. Sci. Math. (2)*, 96:59–64, 1972.
- [12] V. Eberhardt, S. Dierolf, and U. Schwanengel. On the product of two (totally) minimal topological groups and the three-space-problem. *Math. Ann.*, 251(2):123–128, 1980.
- [13] R. Engelking. *General topology*, volume 6 of *Sigma Series in Pure Mathematics*. Heldermann Verlag, Berlin, second edition, 1989. Translated from the Polish by the author.
- [14] E. Hewitt and K. A. Ross. *Abstract harmonic analysis*. Academic Press Inc., Publishers, New York, 1963.
- [15] K. H. Hofmann and S. A. Morris. *The Lie theory of connected pro-Lie groups*, volume 2 of *EMS Tracts in Mathematics*. European Mathematical Society (EMS), Zürich, 2007. A structure theory for pro-Lie algebras, pro-Lie groups, and connected locally compact groups.
- [16] G. Lukács. *Compact-like topological groups*, volume 31 of *Research and exposition in Mathematics*. Heldermann Verlag, Berlin, 2009.
- [17] M. G. Megrelishvili. *G*-minimal topological groups. In *Abelian groups, module theory, and topology (Padua, 1997)*, volume 201 of *Lecture Notes in Pure and Appl. Math.*, pages 289–299. Dekker, New York, 1998.
- [18] P. S. Mostert. Sections in principal fibre spaces. *Duke Math. J.*, 23:57–71, 1956.

- [19] I. R. Prodanov. Precompact minimal group topologies and p -adic numbers. *Annuaire Univ. Sofia Fac. Math.*, 66:249–266 (1974), 1971/72.
- [20] I. R. Prodanov. Some minimal group topologies are precompact. *Math. Ann.*, 227(2):117–125, 1977.
- [21] I. R. Prodanov and L. N. Stojanov. Every minimal abelian group is precompact. *C. R. Acad. Bulgare Sci.*, 37(1):23–26, 1984.
- [22] I. R. Prodanov and L. N. Stojanov. Minimal group topologies. In *Topology, theory and applications (Eger, 1983)*, volume 41 of *Colloq. Math. Soc. János Bolyai*, pages 493–508. North-Holland, Amsterdam, 1985.
- [23] R. M. Stephenson, Jr. Minimal topological groups. *Math. Ann.*, 192:193–195, 1971.
- [24] R. M. Stephenson, Jr. Some unsolved problems concerning P -minimal and P -closed spaces. In *Topology (Proc. Ninth Annual Spring Conf., Memphis State Univ., Memphis, Tenn., 1975)*, pages 249–257. Lecture Notes in Pure and Appl. Math., Vol. 24. Dekker, New York, 1976.
- [25] L. N. Stojanov. On products of minimal and totally minimal groups. In *Proc. 11th Conf. of the Union of Bulg. Math., Sunny Beach 1982*, pages 287–294, 1982.
- [26] M. G. Tkačenko. On dimension of locally pseudocompact groups and their quotients. *Comment. Math. Univ. Carolin.*, 31(1):159–166, 1990.
- [27] T. Yamanoshita. On the dimension of homogeneous spaces. *J. Math. Soc. Japan*, 6:151–159, 1954.

Department of Mathematics and Computer Science
University of Udine
Via delle Scienze, 208 – Loc. Rizzi, 33100 Udine
Italy

Halifax, Nova Scotia
Canada

email: dikranja@dimi.uniud.it

email: lukacs@topgroups.ca